

**bs-11235R****[ Primary Antibody ]****GRK6 Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

|   |                      |  |
|---|----------------------|--|
| <b>Host:</b> Rabbit   | <b>Isotype:</b> IgG  | <b>Applications:</b> <b>IHC-P</b> (1:100-500)<br><b>IHC-F</b> (1:100-500)<br><b>IF</b> (1:100-500)<br><b>ICC/IF</b> (1:100-500)<br><b>ELISA</b> (1:5000-10000)<br><br><b>Reactivity:</b> (predicted: Human, Mouse, Rat, Rabbit, Pig, Cow, Dog)<br><br><b>Predicted MW.:</b> 66 kDa<br><br><b>Subcellular Location:</b> Cell membrane |
| <b>Clonality:</b> Polyclonal  |                      |  |
| <b>GeneID:</b> 2870   | <b>SWISS:</b> P43250 |  |
| <b>Target:</b> GRK6   |                      |  |
| <b>Immunogen:</b> KLH conjugated synthetic peptide derived from human GRK6: 181-280/576.  |                      |  |
| <b>Purification:</b> affinity purified by Protein A   |                      |  |
| <b>Concentration:</b> 1mg/ml  |                      |  |
| <b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.<br>Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.  |                      |  |
| <b>Background:</b> Heterotrimeric G protein-mediated signal transduction is a dynamically regulated process with the intensity of signal decreasing over time despite the continued presence of the agonist. This phenomenon, referred to as agonist-mediated desensitization, involves phosphorylation of the receptor by two classes of enzymes. The first class is comprised of the second messenger-regulated kinases, such as c-AMP dependent protein kinase A and protein kinase C. The second class includes the G protein-coupled receptor kinases (GRKs). At least seven members of the GRK family have been identified. These include rhodopsin kinase (GRK 1), two forms of Beta-adrenergic receptor kinase: GRK 2 (Beta ARK, Beta ARK1) and GRK 3 (Beta ARK2), IT-11 (GRK 4), GRK 5, GRK 6 and GRK 7. Phosphorylation of receptors by GRKs appears to be strictly dependent on the receptor being in its agonist-activated state. |                      |  |