bs-11258R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Rab11A Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 8766 SWISS: P62491

Target: Rab11A

Immunogen: KLH conjugated synthetic peptide derived from human Rab11A:

141-216/216.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Rab proteins are low-molecular-weight GTP-binding proteins that form the largest branch of the Ras superfamily of GTPases. Located on the cytoplasmic face of organelles and vesicles, rab proteins are involved in intracellular membrane fusion reactions. Three membrane proteins, synaptosomal associated protein of 25 kDa (SNAP-25), synaptobrevin, and syntaxin, form the core of a ubiquitous membrane fusion machine that interacts with the soluble proteins N-ethylmaleimide-sensitive factor (NSF) and a-SNAP. Rab proteins, in coordination with the core fusion machinery and Munc-18, help to mediate vesicle docking and fusion. There are over 40 Rab proteins in mammals. Rab11a and Rab11b are known markers for protein trafficking, sorting, and recycling in the endosomal pathway. The Rab11 proteins are enriched in recycling endosomes and the trans-Golgi network, where they regulate membrane recycling back to the plasma membrane. Rab11a is ubiquitously expressed, while Rab11b is expressed mainly in the heart and brain.

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) **IF** (1:100-500)


Reactivity: Human (predicted: Mouse,

Rat, Cow)

Predicted 24 kDa MW.:

Subcellular Location: Cell membrane ,Cytoplasm

VALIDATION IMAGES -



Sample: Lane 1: Recombinant human Rab11 protein, N-His(bs-42325P) Primary: Anti-Rab11A (bs-11258R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 24 kDa Observed hand size: 26 kDa



Paraformaldehyde-fixed, paraffin embedded (human heart); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (Rab11A) Polyclonal Antibody, Unconjugated (bs-11258R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

• [IF=12.5] Zhenyu Wang, et al. Morphology– and adhesion–dual biomimetic nanovaccine boosts antigen crosspresentation through subcellular transport regulation. SCI ADV. 2025 Jul;11(31) IF; Mouse. 40737412