bs-19984R

[Primary Antibody]

BIOSS ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

PRDM13 Rabbit pAb

GeneID: 59336 **SWISS:** Q9H4Q3

Target: PRDM13

Immunogen: KLH conjugated synthetic peptide derived from human PRDM13:

551-650/707.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: PRDM13 may be involved in transcriptional regulation.

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)

Reactivity: Mouse, Hamster


(predicted: Human, Rat, Rabbit, Pig, Sheep, Cow,

Dog)


Predicted MW.: 74 kDa

Subcellular Nucleus

- VALIDATION IMAGES -



Sample: CHO(Hamster) Cell Lysate at 30 ug Primary: Anti-PRDM13 (bs-19984R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 74 kD Observed band size: 74 kD



Paraformaldehyde-fixed, paraffin embedded (mouse testis tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PRDM13) Polyclonal Antibody, Unconjugated (bs-19984R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.