

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## S-tag Rabbit pAb

Catalog Number: bs-33017R

Target Protein: S-tag Concentration: 1mg/ml

> Form: Liquid Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:1000-5000)

Reactivity: Species independent

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: S-tag is the name of an oligopeptide derived from pancreatic ribonuclease A (RNase A). If

RNase A is digested with subtilisin, a single peptide bond is cleaved, but the resulting two

products remain weakly bound to each other and the protein, called ribonuclease S,

remains active although each of the two products alone shows no enzymatic activity. The Nterminus of the original RNase A, also called S-peptide, consists of 20 amino acid residues, of


which only the first 15 are required for ribonuclease activity. This 15 amino acids long

peptide is called S15 or S-tag.

The amino acid sequence of the S-tag is: Lys-Glu-Thr-Ala-Ala-Lys-Phe-Glu-Arg-Gln-His-Met-Asp-Ser. It is believed that the peptide with its abundance of charged and polar residues could improve solubility of proteins it is attached to [citation needed]. Moreover, the peptide alone is thought not to fold into a distinct structure. On DNA-level the S-tag can be attached to the N- or C-terminus of any protein. After gene expression, such a tagged protein can be

detected by commercially available antibodies.

## **VALIDATION IMAGES**



Sample: Lane1: S-Tagged Fusion Protein Overexpression E.coli Lysate at 2ug Lane2: S-Tagged Fusion Protein Overexpression E.coli Lysate at 2ug Primary: Anti-S-tag (bs-33017R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 41 kD Observed band size: 45 kD

## PRODUCT SPECIFIC PUBLICATIONS

[IF=5.486] Zhangji Dong. et al. Histone acetyltransferase KAT2A modulates neural stem cell differentiation and proliferation by inducing degradation of the transcription factor PAX6. J BIOL CHEM. 2023 Feb;:103020 IP; Human . 36791914