### bs-33017R

## [ Primary Antibody ]

# Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# S-tag Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Target: S-tag

Purification: affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** S-tag is the name of an oligopeptide derived from pancreatic

ribonuclease A (RNase A). If RNase A is digested with subtilisin, a single peptide bond is cleaved, but the resulting two products remain weakly bound to each other and the protein, called ribonuclease S, remains active although each of the two products alone shows no enzymatic activity. The N-terminus of the original RNase A, also called S-peptide, consists of 20 amino acid residues, of which only the first 15 are required for ribonuclease activity. This


15 amino acids long peptide is called S15 or S-tag.
The amino acid sequence of the S-tag is: Lys-Glu-Thr-Ala-Ala-Ala-Lys-Phe-Glu-Arg-Gln-His-Met-Asp-Ser. It is believed that the peptide with its abundance of charged and polar residues could improve solubility of proteins it is attached to [citation needed]. Moreover, the peptide alone is thought not to fold into a distinct structure. On DNA-level the S-tag can be attached to the N- or C-terminus of any protein. After gene expression, such a tagged protein can be detected by commercially available antibodies.

Applications: WB (1:1000-5000)

IHC-P (1:200-1000) IHC-F (1:200-1000) IF (1:200-1000) Flow-Cyt (1ug/Test) ICC/IF (1:100-500) ELISA (1:1000-5000)

Reactivity: Species independent

#### - VALIDATION IMAGES -



Sample: Lane1: S-Tagged Fusion Protein
Overexpression E.coli Lysate at 2ug Lane2: STagged Fusion Protein Overexpression E.coli
Lysate at 2ug Primary: Anti-S-tag (bs-33017R) at
1/1000 dilution Secondary: IRDye800CW Goat
Anti-Rabbit IgG at 1/20000 dilution Predicted
band size: 41 kD Observed band size: 45 kD

#### - SELECTED CITATIONS -

• [IF=5.486] Zhangji Dong. et al. Histone acetyltransferase KAT2A modulates neural stem cell differentiation and proliferation by inducing degradation of the transcription factor PAX6. J BIOL CHEM. 2023 Feb;:103020 IP; Human. 36791914