

bs-18527R**[Primary Antibody]****ZFP161 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)
Clonality: Polyclonal		
GeneID: 7541	SWISS: O43829	
Target: ZFP161		Reactivity: Rat (predicted: Human, Mouse, Rabbit, Cow, Dog, Horse)
Immunogen: KLH conjugated synthetic peptide derived from human ZFP161: 181-280/449.		
Purification: affinity purified by Protein A		Predicted MW.: 51 kDa
Concentration: 1mg/ml		Subcellular Location: Nucleus
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: ZFP161 is a 449 amino acid member of the Krüppel C2H2-type zinc-finger protein family. Localized to distinct regions within the nucleus, ZFP161 functions both as a transcriptional repressor of Thymidine Kinase (TK) and c-Myc promoters and as a transcriptional activator of the dopamine transporter (DAT) promoter. ZFP161 contains one BTB (POZ) domain and five C2H2-type zinc fingers which direct its specific localization and nucleic acid binding, respectively. Defects in the gene encoding ZFP161 may be associated with holoprosencephaly type 4 (HPE4), a structural anomaly of the brain characterized by a flattened nasal tip with no visible septum, hypotelorism, lack of nasal bridge, ptosis of the left upper eyelid and, in some cases, infant death.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (Rat ovary); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZFP161) Polyclonal Antibody, Unconjugated (bs-18527R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.