

**bsm-52614R****[ Primary Antibody ]**

**Bioss**  
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

## ACE2 Recombinant Rabbit mAb

### DATASHEET

**Host:** Rabbit**Clonality:** Recombinant**GeneID:** 59272**Target:** ACE2**Immunogen:** A synthesized peptide derived from human ACE2: 200-226. < Extracellular >**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Store at -20°C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4°C.

**Background:** The protein encoded by this gene belongs to the angiotensin-converting enzyme family of dipeptidyl carboxydipeptidases and has considerable homology to human angiotensin 1 converting enzyme. This secreted protein catalyzes the cleavage of angiotensin I into angiotensin 1-9, and angiotensin II into the vasodilator angiotensin 1-7. The organ- and cell-specific expression of this gene suggests that it may play a role in the regulation of cardiovascular and renal function, as well as fertility. In addition, the encoded protein is a functional receptor for the spike glycoprotein of the human coronaviruses SARS and HCoV-NL63. [provided by RefSeq, Jul 2008]


**Isotype:** IgG**CloneNo.:** 3F1**SWISS:** Q9BYF1**Applications:** WB (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**ICC/IF** (1:50-200)**Reactivity:** Human, Mouse, Rat**Predicted MW.:** 92 kDa

**Subcellular Location:** Cytoplasm ,Cell projection  
Membrane ,Secreted ,Cell membrane


### VALIDATION IMAGES




Sample: Lane 1: Recombinant human ACE2 protein, His & Avi (HEK293)( bs-46001P) Primary: Anti-ACE2 (bsm-52614R) at 1/1000 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 92 kDa  
Observed band size: 105 kDa




25 ug total protein per lane of various lysates (see on figure) probed with ACE2 monoclonal antibody, unconjugated (bsm-52614R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.




Paraformaldehyde-fixed, paraffin embedded (rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ACE2) Monoclonal Antibody, Unconjugated (bsm-52614R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining



Paraformaldehyde-fixed, paraffin embedded



Paraformaldehyde-fixed, paraffin embedded



Paraformaldehyde-fixed, paraffin embedded

**Important Note:** This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Human Testicles; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with ACE2 Monoclonal Antibody, Unconjugated (bsm-52614R) at 1:200 overnight at 4°C, followed by conjugation to the SP Kit (Rabbit, SP-0023) and DAB (C-0010) staining.

Human Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with ACE2 Monoclonal Antibody, Unconjugated (bsm-52614R) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-0295G-BF488), DAPI (blue, C02-04002) was used to stain the cell nuclei.

Mouse Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with ACE2 Monoclonal Antibody, Unconjugated (bsm-52614R) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-0295G-BF488), DAPI (blue, C02-04002) was used to stain the cell nuclei.

---

## — SELECTED CITATIONS —

- **[IF=5.776]** Endika Prieto-Fernández. et al. Hypoxia reduces cell attachment of SARS-CoV-2 spike protein by modulating the expression of ACE2, neuropilin-1, syndecan-1 and cellular heparan sulfate. *Emerg Microbes Infect.* 2021;10(1):1065-1076  
WB ;Human. 34013835