

bs-14657R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

EXOC1 Rabbit pAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500)
GeneID: 55763	SWISS: Q9NV70	IF (1:100-500)
Target: EXOC1		Reactivity: Rat (predicted: Human, Mouse, Pig, Sheep, Cow, Horse)
Immunogen: KLH conjugated synthetic peptide derived from human EXOC1: 701-800/894.		Predicted MW.: 102 kDa
Purification: affinity purified by Protein A		Subcellular Location: Cell membrane ,Cytoplasm
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Exocytosis is crucial in membrane trafficking and it mediates hormone and neurotransmitter secretion out of the cell, as well as the incorporation of membrane proteins and lipids to the plasma membrane. It is crucial for cell-cell communication, cell growth and cell polarity. The exocyst complex is a multi-protein complex that consists of Sec3, Sec5, Sec6, Sec8, Sec10, Sec15, Exo70 and Exo84, and is essential for targeting exocytic vesicles to specific docking sites on the plasma membrane. Exo70, also known as EXOC7 (exocyst complex component 7), EXOC1 or 2-5-3p, is a 735 amino acid peripheral membrane protein that is a component of the exocyst complex. Localized to the cytoplasm and the cell membrane, Exo70 plays an essential role in the docking of exocystic vesicles to target sites on the plasma membrane and, specifically, may be involved in insulin-induced protein docking within the cell. Four isoforms of Exo70 are expressed due to alternative splicing events.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EXOC1) Polyclonal Antibody, Unconjugated (bs-14657R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.