bs-8158R

[Primary Antibody]

HCV NS5A transactivated protein 8 Rabbit pAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 53339 SWISS: Q9H0C5

Target: HCV NS5A transactivated protein 8

Immunogen: KLH conjugated synthetic peptide derived from human

BTBD1/HCV NS5A transactivated protein 8: 401-482/482.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: BTBD1 contains a proline-rich region and a BTB/POZ domain

which are typically involved in protein-protein interactions. The Cterminus of BTBD1 binds topoisomerase I and BTBD1 also interacts with TRIM5 isoform Delta. BTBD1 localizes to cytoplasmic bodies

and is essential for myogenic differentiation.

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) **IF** (1:100-500) **ELISA** (1:5000-10000)

Reactivity: (predicted: Human, Mouse,

Rat, Pig, Sheep, Cow, Dog,

Horse)

Predicted MW.: 53 kDa

Subcellular Cytoplasm