

**bs-14663R**

**[ Primary Antibody ]**

## EXOC6B Rabbit pAb

**Bioss**  
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800


### — DATASHEET —

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> <b>WB</b> (1:500-2000) <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500)  <b>Reactivity:</b> Human, Rat (predicted: Mouse, Rabbit, Pig, Sheep, Cow, Dog, Horse)  <b>Predicted MW.:</b> 94 kDa  <b>Subcellular Location:</b> Cytoplasm
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 23233	<b>SWISS:</b> Q9Y2D4	
<b>Target:</b> EXOC6B		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human EXOC6B: 721-811/811.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> In yeast and rat, Sec15 is part of a multiprotein complex that is required for targeted exocytosis.[supplied by OMIM, Apr 2004]		


### — VALIDATION IMAGES —



Sample: HeLa (Human) Cell Lysate at 40 ug  
Primary: Anti-EXOC6B(bs-14663R) at 1/300  
dilution Secondary: IRDye800CW Goat Anti-  
RabbitIgG at 1/20000 dilution Predicted band  
size: 94kD Observed band size: 94kD



Paraformaldehyde-fixed, paraffin embedded  
(Rat small intestine); Antigen retrieval by boiling  
in sodium citrate buffer (pH6.0) for 15min; Block  
endogenous peroxidase by 3% hydrogen  
peroxide for 20 minutes; Blocking buffer (normal  
goat serum) at 37°C for 30min; Antibody  
incubation with (EXOC6B) Polyclonal Antibody,  
Unconjugated (bs-14663R) at 1:500 overnight at  
4°C, followed by a conjugated secondary  
(sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded  
(Rat stomach); Antigen retrieval by boiling in  
sodium citrate buffer (pH6.0) for 15min; Block  
endogenous peroxidase by 3% hydrogen  
peroxide for 20 minutes; Blocking buffer (normal  
goat serum) at 37°C for 30min; Antibody  
incubation with (EXOC6B) Polyclonal Antibody,  
Unconjugated (bs-14663R) at 1:500 overnight at  
4°C, followed by a conjugated secondary  
(sp-0023) for 20 minutes and DAB staining.