

EXOC6B Rabbit pAb

Catalog Number: bs-14663R

Target Protein: EXOC6B

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Rat (predicted: Mouse, Rabbit, Pig, Sheep, Cow, Dog, Horse)

Predicted MW: 94 kDa

Subcellular: Cytoplasm

Locations:

Entrez Gene: 23233

Swiss Prot: Q9Y2D4

Source: KLH conjugated synthetic peptide derived from human EXOC6B: 721-811/811.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: In yeast and rat, Sec15 is part of a multiprotein complex that is required for targeted exocytosis.[supplied by OMIM, Apr 2004]


VALIDATION IMAGES



Sample: HeLa (Human) Cell Lysate at 40 ug Primary: Anti-EXOC6B(bs-14663R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 94kD Observed band size: 94kD



Paraformaldehyde-fixed, paraffin embedded (Rat stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EXOC6B) Polyclonal Antibody, Unconjugated (bs-14663R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat small intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EXOC6B) Polyclonal Antibody, Unconjugated (bs-14663R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.