

bs-14667R**[Primary Antibody]****EXOSC4 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit Clonality: Polyclonal GeneID: 54512 Target: EXOSC4 Immunogen: KLH conjugated synthetic peptide derived from human EXOSC4: 51-150/245. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: The exosome is a multisubunit complex composed of several highly conserved subunits, some of which are 3' to 5' exoribonucleases. The complex is involved in a variety of cellular processes and is responsible for degrading unstable mRNAs that contain AU-rich (ARE) elements in their untranslated 3' region. EXOSC4 (exosome component 4), also known as SKI6, RRP41 or p12A, is a 245 amino acid protein that localizes to both the nucleus and the cytoplasm and shares 96% sequence identity with its mouse counterpart. Functioning as a component of the exosome complex, EXOSC4 exhibits 3'-5' exonuclease activity and is required for the 3'-processing of 7S pre-rRNA to mature 5.8S rRNA. The gene encoding EXOSC4 maps to human chromosome 8, which consists of nearly 146 million base pairs, houses more than 800 genes and is associated with a variety of diseases and malignancies.	Isotype: IgG SWISS: Q9NPD3 Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Mouse (predicted: Human, Rat, Pig, Sheep, Cow, Dog, Horse) Predicted MW.: 26 kDa Subcellular Location: Cytoplasm ,Nucleus
---	---

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EXOSC4) Polyclonal Antibody, Unconjugated (bs-14667R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.